

NO-TILL WHEAT INCREASES YIELDS OF CORN AND SOYBEANS IN THE CROPPING ROTATION

L. Murdock, J. Herbek, J. Martin, J. James, and D. Call
Plant & Soil Sciences Department
University of Kentucky, Princeton, KY 42445
PH: (270) 365-7541, Ext. 207; Email: lmurdock@uky.edu

OBJECTIVE

The objective of this experiment was to verify the effects of no-till wheat and tilled wheat on the subsequent yield of soybeans and corn planted after wheat in a wheat, double-cropped soybean and corn rotation and measure differences in fertility and physical effects on the soil on a long-term basis.

METHODS

The experiment began in the fall of 1992 at Princeton, Ky on a Huntington silt loam soil that is moderately well drained. Wheat was planted no-till and with tillage and the tillage plots were chisel plowed and disked twice. The plots were 10 ft x 30 ft. The trial was soil tested each year and fertilizer and lime applied according to University of Kentucky recommendations. N was sidedressed on corn at 150 lb/ac. Soybeans are planted no-till immediately after wheat harvest and no-till corn is planted the following year and wheat (tilled and no-tilled) is again planted after corn harvest.

RESULTS

Yields of Succeeding Crops

The data (Table 1) indicates that both no-till corn and no-till soybeans tend to yield more (5.0% for soybeans and 3.5% for corn) where the wheat is planted no-till. Over the 18 years of the trial, either corn or soybean yields have been statistically higher 8 of the 18 years (about 50% of the years) when planted after no-till wheat compared to tilled wheat. The actual yield increase has been 2.1 bu/ac. for soybeans and 6.8 bu/ac. for the corn. With prices for corn at \$5.00/bu and soybean at \$10.00/bu, this

would mean a \$55.00 increased return over the two year rotation for planting no-till wheat.

These yield differences indicate that changes between the two systems have taken place with time and the changes favor the system which is completely no-tillage including the no-till wheat. Research indicates that the reason for the difference is due to residue cover, soil moisture, soil physical changes and more importantly a change in pore size distribution. There are more medium sized pores in the upper few inches of the soil that hold more plant available water.

Soil Changes

There is no difference in the soil density between the systems. This indicates that there was no compaction of significance in either system. The soil strength, as indicated by penetrometer measurements, was higher in the exclusively no-tillage system. This and other soil measurements indicate that the soil structure has changed and has larger aggregates and more medium sized pores than the system that is tilled every second year for wheat planting. An advantage for this change in pore size was seen in 2004. The wet soil conditions after wheat harvest for planting double-cropped soybeans were different for the two tillage systems. The soil with no-till wheat had better drainage and better planting conditions than the tilled wheat treatment. This resulted in 30% more plants than the tilled wheat treatment.

Soil moisture measurements taken during the latter part of the growing season are higher in

the true no-till system about half of the years and about the same as tilled the other years. When the soil moisture is higher, corn and soybean yields are usually higher in the no-till system. In the years when there is no difference in soil moisture, the yields are similar. The no-tilled soil can hold more plant available water but the rain must fall at the proper time for the advantage to express itself. In some years, it does and others it does not.

SUMMARY AND CONCLUSIONS

A true no-tillage system seems to have a favorable effect on the corn and soybean crops grown in rotation with the no-till wheat. When no-till wheat was grown, the average yields of no-till corn and soybeans had 3.5 and 5.0% greater yields, respectively, than when these crops were grown after tilled wheat. This increase in yield results in about a \$55/ac. increase in profits for the 2 year rotation of wheat, double-cropped soybeans and corn. The soil changes include larger aggregates and more medium pores which result in more plant available moisture for these crops.

Table 1. Effect on Tilled and No-Tilled Wheat on Yields of No-Tilled Corn and Soybeans in a Cropping Rotation				
----- Yield (bu/ac) -----				
	Soybean		Corn	
Number of Continuous Years	N.T. Wheat	Tilled Wheat	N.T. Wheat	Tilled Wheat
18	43.5	41.4	196.2	189.4